

Germline ESR2 mutation predisposes to medullary thyroid carcinoma and causes up-regulation of RET expression

Smith, Joel; Read, Martin; Hoffman, Jon; Brown, Rachel; Bradshaw, Beth; Campbell, Christopher; Cole, Trevor; Dieguez Navas, Johanna; Eatock, Fiona; Gundara, Justin; Lian, Eric; McMullan, Dominic; MacDonald, Fiona; Morgan, Neil; Mulligan, Lois; Morrison, Patrick J.; Robledo, Mercedes; Simpson, Michael; Smith, Vicki; Stewart, Sue

DOI:

[10.1093/hmg/ddw057](https://doi.org/10.1093/hmg/ddw057)

Document Version

Peer reviewed version

Citation for published version (Harvard):

Smith, J, Read, M, Hoffman, J, Brown, R, Bradshaw, B, Campbell, C, Cole, T, Dieguez Navas, J, Eatock, F, Gundara, J, Lian, E, McMullan, D, MacDonald, F, Morgan, N, Mulligan, L, Morrison, PJ, Robledo, M, Simpson, M, Smith, V, Stewart, S, Trembath, R, Sidhu, S, Togneri, F, Vialard, L, Wake, N, Wallis, Y, Watkinson, J, Maher, E, McCabe, C & Woodward, E 2016, 'Germline ESR2 mutation predisposes to medullary thyroid carcinoma and causes up-regulation of RET expression', *Human Molecular Genetics*, vol. 25, no. 9, pp. 1836-1845.
<https://doi.org/10.1093/hmg/ddw057>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:

Authors may upload their accepted manuscript PDF to an institutional and/or centrally organized repository, provided that public availability is delayed until 12 months after first online publication in the journal. Note: The article was published online on 3rd March 2016.

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Germline ESR2 Mutation Predisposes to Medullary Thyroid Carcinoma and Causes Up-Regulation of RET Expression

Joel Smith^{#1}, Martin Read^{#2}, Jon Hoffman³, Rachel Brown⁴, Beth Bradshaw⁵, Christopher Campbell^{5, 15}, Trevor Cole³, Johanna Dieguez Navas⁶, Fiona Eatock⁷, Justin S. Gundara⁸, Eric Lian⁹, Dom McMullan⁵, Neil V. Morgan², Lois Mulligan⁹, Patrick J Morrison¹⁰, Mercedes Robledo¹¹, Michael A. Simpson¹², Vicki Smith², Sue Stewart³, Richard C Trembath¹³, Stan Sidhu⁸, Fiona S Togneri⁵, Naomi C Wake¹, Yvonne Wallis⁵, John C Watkinson⁴, Eamonn R Maher^{2, 14}, Chris Mc Cabe², and Emma R Woodward^{* 1, 3, 15}

[#] Joint first authors

1. Centre for Rare Diseases and Personalised Medicine, University of Birmingham, Birmingham B15 2TT, UK

2. School of Clinical and Experimental Medicine, University of Birmingham, B15 2TT, UK

3. West Midlands Regional Genetics Service, Birmingham Women's Hospital, Birmingham B15 2TG, UK

4. Queen Elizabeth Hospital, Queen Elizabeth Medical Centre, Birmingham B15 2TH, UK

5. West Midlands Regional Genetics Laboratory, Birmingham Women's Hospital, Birmingham B15 2TG, UK

6. Human Biomaterials Resource Centre, College of Medical and Dental Sciences, University of Birmingham, Vincent Drive, Edgbaston B15 2TT, UK

7. Department of Endocrine Surgery, Belfast Health and Social Care Trust, Royal Victoria Hospital, Belfast, Northern Ireland, UK

8. Cancer Genetics, Level 9, Kolling Building and Endocrine Surgical Unit, Royal North Shore Hospital, University of Sydney, Pacific Highway, St. Leonards, NSW, 2065, Australia

9. Division of Cancer Biology and Genetics, Cancer Research Institute, Queen's University, Canada

10. Centre for Cancer Research and Cell Biology, Queen's University of Belfast, 97 Lisburn Road, Belfast BT9 7AE

11. Hereditary Endocrine Cancer Group, Spanish National Cancer Research Centre (CNIO), Madrid, Spain and Centro de Investigación Biomédica en Red de Enfermedades Raras, Madrid, Spain.

12. Division of Genetics and Molecular Medicine, King's College London School of Medicine, Guy's Hospital, London, UK

13. Queen Mary University of London, Barts and The London School of Medicine and Dentistry, London, UK

14. Department of Medical Genetics, University of Cambridge and NIHR Cambridge Biomedical Research Centre, Cambridge, UK

15. Manchester Centre for Genomic Medicine, Central Manchester University Hospitals NHS Foundation Trust, Manchester Academic Health Sciences Centre (MAHSC), Manchester M13 9WL, UK

* corresponding author to whom correspondence should be addressed

Dr. Emma Woodward

Manchester Centre for Genomic Medicine

St Mary's Hospital

Oxford Road

Manchester, M13 9WL

United Kingdom

Telephone: +44 161 276 6506

Fax: +44 161 276 6145

Email: emma.woodward@cmft.nhs.uk

Abstract

Familial medullary thyroid cancer (MTC) and its precursor, C cell hyperplasia (CCH), is associated with germline *RET* mutations causing multiple endocrine neoplasia type 2. However, some rare families with apparent MTC/CCH predisposition do not have a detectable *RET* mutation. To identify novel MTC/CCH predisposition genes we undertook exome resequencing studies in a family with apparent predisposition to MTC/CCH and no identifiable *RET* mutation. We identified a novel *ESR2* frameshift mutation, c.948delT, which segregated with histological diagnosis following thyroid surgery in family members and demonstrated loss of *ESR2* encoded ER β expression in the MTC tumour. ER α and ER β form heterodimers binding DNA at specific estrogen response elements (ERE) to regulate gene transcription. ER β represses ER α mediated activation of the ERE and the *RET* promoter contains three ERE. *In vitro*, we showed that *ESR2* c.948delT results in unopposed ER α mediated increased cellular proliferation, activation of the ERE and increased RET expression. *In vivo*, immunostaining of CCH and MTC using an anti-RET antibody demonstrated increased RET expression. Together these findings identify germline *ESR2* mutation as a novel cause of familial MTC/CCH and provide important insights into a novel mechanism causing increased RET expression in tumourigenesis.

Introduction

Thyroid cancer is the most common endocrine malignancy. Although medullary thyroid cancer (MTC), arising from the calcitonin secreting para-follicular cells, only accounts for ~5% of thyroid cancers (1), nearly 50% of patients present with stage III or IV disease and only 21% of patients presenting with stage IV disease survive 10 years (2).

In ~25% of MTC there is a germline *RET* mutation (3) predisposing to multiple endocrine neoplasia type 2 (MEN2), a dominantly inherited endocrine tumour predisposition syndrome characterised by predisposition to the development of young onset MTC and often primary C cell hyperplasia (CCH), a precursor to MTC (4). Other features may also occur such as pheochromocytoma, primary hyperparathyroidism (HPT) and, rarely, developmental abnormalities, for example, a marfanoid habitus and ganglioneuromatosis of the mouth and gut (4).

The majority of *RET* mutations predisposing to MEN2 result in single amino acid substitutions affecting key residues in the extra-cellular and kinase domains of the RET receptor (5,6) causing inappropriate constitutive RET activation. Detection of a germline *RET* mutation, enables pre-symptomatic interventions such as prophylactic thyroidectomy to be offered to at-risk gene carriers (6). Thus all individuals presenting with MTC or primary CCH should be offered germline *RET* testing (7); however, some families with an apparent predisposition to MTC/CCH do not harbour a germline *RET* gene alteration (8,9) suggesting that further predisposing gene alterations remain to be identified. We investigated a kindred with non-*RET* MTC/CCH and detected a novel constitutional frameshift mutation in *ESR2* encoding the beta subunit of the oestrogen receptor, ER β (10).

Results

Case Report

The index case presented age 22 years with MTC (Supplementary Material, Fig. S1). Although no constitutional mutation of exons 10, 11, or 13-16 of *RET* was detected, in view of the young age of diagnosis his monozygotic twin brother was offered follow-up and underwent thyroidectomy age 33 years because of abnormal pentagastrin stimulation testing; histological review showed the presence of CCH. Prophylactic thyroid surgery was then undertaken in the offspring of both brothers (individuals III:1-5); individuals III:2 and III:3 were found to have CCH and individuals III:1, III:4 and III:5 to have normal thyroid tissue. Pentagastrin stimulation testing was subsequently offered to the wider family and individual I:1 noted to have mildly abnormal results. In the index case, further constitutional molecular genetic testing did not detect any mutations in the remaining exons of *RET*, karyotype analysis was normal and array CGH analysis did not demonstrate any clinically significant CNVs (data not shown). We therefore hypothesised a novel gene alteration was predisposing to MTC/CCH in the family and undertook exome sequencing in two affected members (II:3 and III:3).

Identification of Germline *ESR2* mutation in Familial MTC/CCH

We filtered the variants identified using the dbSNP (11), NHLBI Exome Variant Server (EVS) (12) and 1000 genomes (13) datasets looking for potentially disrupting variants present in both samples, prioritising frameshift and nonsense changes. We did not identify any deleterious alterations in the genes known to be mutated in hereditary cancer predisposition (14, Supplementary Material, Table S1) but did identify a novel frameshift alteration of *ESR2* (c.948delT; p.Gly318Alafs*22) present in both affected individuals but not in the NHLBI EVS nor in 2577 individuals from the 1000 genome project (12,13).

Constitutional *ESR2* sequencing in other family members demonstrated the c.948delT variant to be present in the two family members found to have CCH (individuals II:1 and III:2) but not in the

three (individuals III:1, III:4, III:5) with normal thyroid histology (Fig. 1A, Supplementary Material, Fig. S1).

Loss of ER β staining in *ESR2* c.948delT associated MTC

To investigate the role of *ESR2* in MTC tumourigenesis we initially undertook immunohistochemistry using an anti-ER β antibody and demonstrated complete loss of nuclear staining in the c.948delT associated MTC (Fig. 1B) with retained staining in the corresponding normal thyroid tissue. However, despite direct sequencing of the complete *ESR2* coding region and LOH studies using closely linked microsatellite markers in archived tumour material, we did not detect a somatic alteration of *ESR2* (data not shown). Whole genome copy number analysis by molecular inversion probe technology did not reveal any pathological CNVs affecting either chromosomes 14 or 10 (harbouring the *ESR2* and *RET* loci respectively) (Supplementary Material, Fig. S2).

No additional constitutional *ESR2* mutations in other cases of apparent MTC predisposition

To ascertain whether constitutional alterations of *ESR2* might be involved in other cases of apparent MTC predisposition, we undertook sequencing of the coding region of *ESR2* in 19 individuals with apparently isolated MTC, eight of whom were diagnosed <40 years of age (mean age at diagnosis 47.5y, range 33-71y, SEM 2.96). The entire coding region of *RET* had been previously sequenced and no alterations detected other than known polymorphisms. We identified a novel germline missense alteration, c.382G>C; p.V128L, in a female who developed an apparently isolated MTC age 36 years (Fig. 2A). This variant was also detected in a pair of siblings, one of whom had a papillary thyroid cancer age 60 years and the other a medullary thyroid cancer age 57 years. We did not detect any constitutional *ESR2* alterations, other than known polymorphisms, in a further three families with familial non-*RET* MTC (data not shown).

Further investigation using archived tumour material from the individual with the apparently isolated MTC and *ESR2* c.382G>C (no blocks were available from the sibling pair) revealed preserved ER β staining (Fig. 2B) and no somatic *ESR2* sequence alterations, LOH or pathological

CNVs of chromosomes 14 or 10 were detected. Although this variant was not present in the NHLBI EVS (12) or 1000 genome project (13) datasets, further analysis using SIFT (15) predicted it to be tolerated, and PolyPhen-2 (16) to be benign; we were therefore unable to exclude that *ESR2* c.382G>C may represent a rare polymorphism and did not evaluate further.

No somatic mutations of *ESR2* detected in sporadic MTC

As hereditary cancer predisposing genes are often implicated in sporadic tumours (17), we investigated whether somatic mutations of *ESR2* also occurred in sporadic MTC. However, direct sequencing of *ESR2* in 15 fresh frozen sporadic MTC did not reveal any alterations other than known polymorphisms (data not shown). Whilst the cosmic (18) and TCGA data sets (19), containing data for over 21,000 tumours, did not have any data regarding *ESR2* in MTC, we did note the presence of somatic *ESR2* mutations in other tumours of neuroectodermal origin, for example glioma and melanoma.

***ESR2* c.948delT is associated with unrestrained ER α driven cell proliferation**

We then sought to understand how this rare novel *ESR2* frameshift alteration might cause tumorigenesis. Initially we determined stability of the *ESR2* c.948delT mutant compared to wild-type (wt) *ESR2*. Transient transfection of HCT116 cells, followed by the addition of the transcription inhibitor, actinomycin, resulted in significantly lower *ESR2* c.948delT mRNA levels over a 24 hour time period compared with wt*ESR2* indicating reduced stability of the mutant mRNA (Supplementary Material, Fig. S3).

ESR2-encoded ER β forms either homo- or, preferentially, hetero-dimers with *ESR1*-encoded ER α to bind DNA at specific estrogen-responsive elements (EREs) within the promoters of target genes to regulate transcription (20). ER α is the more potent activator and ER β can repress ER α ; thus transcriptional activity is determined by the relative ER α /ER β proportion (21-23).

We next investigated whether *ESR2* c.948delT altered cellular growth. In MCF-7 cells, *ESR2* c.948delT failed to restrain the ER α -driven proliferation of MCF-7 cells in response to either 17 β -

estradiol (E2) or the ER α selective agonist, PPT. In contrast, wtESR2 was capable of inhibiting cell proliferation by up to 30% in both E2 and PPT treated MCF7 cells over-expressing ER α (Fig. 3).

Loss of ER β function causes increased estrogen-responsive element (ERE) activity and RET expression

In both HCT116 and MCF-7 cells, in response to either E2 or PPT agonist, wtESR1 activated the ERE, as indicated by increased luciferase expression, and this was significantly restrained in the presence of wtESR2. However, co-transfection of wtESR1 with ESR2 c.948delT restored expression to a level similar to that seen with wtESR1 alone indicating that the ESR2 c.948delT mutant has lost the ability to restrain ER α -mediated activation of the ERE (Fig. 4).

We then investigated how inappropriate activation of the ERE by c.948delT might cause MTC tumourigenesis. Inappropriate up-regulation of RET activity causing MTC is well established (4) and we noted that the RET promoter (-34 to -314) contains three ERE (24). Furthermore, in *in vitro* studies, RET has been established as an ER target gene (25-27). We therefore investigated whether loss of ER β function might lead to up-regulation of RET expression.

As MCF7 cells have intrinsic ER activity, we used HCT116 cells to investigate the effect of ESR2 c.948delT on RET expression. Co-transfection of wtESR1 with wtESR2 in HCT116 cells, in the presence of either E2 or PPT, resulted in decreased RET expression as compared with wtESR1 alone; however, over-expression of wtESR1 and ESR2 c.948delT, restored RET expression to a level similar to that seen with wtESR1 alone (Fig. 5A, B).

In vivo, RET immunostaining using archived sections from the ESR2 c.948delT associated MTC (individual II:3) and analysis of archived stained slides showing CCH (individual II:1) demonstrated increased RET staining (Fig. 5C). In particular, the intensity of RET expression in the MTC was stronger than in corresponding normal thyroid tissue and similar to that in an MEN2-associated MTC with a known constitutional RET mutation (Fig. 5C). Analysis of RET exons 10, 11 and 13-16 in the ESR2 c.948delT associated MTC did not detect a somatic mutation to account for the increased expression (data not shown).

Discussion

By targeted capture array-based exome sequencing we detected a novel loss of function *ESR2* mutation in a family with predisposition to MTC/CCH. As the interpretation of novel rare variants identified through next generation sequencing approaches in isolated families is challenging, we sought functional evidence to establish pathogenicity.

Initially we demonstrated tumour specific loss of ER β expression in the c.948delT associated MTC which we considered indicative of loss of function as ER β is known to be expressed by human para-follicular C cells (28). Whilst *ESR2* knockout mice display a phenotype more in keeping with a reduced fertility phenotype in females rather than one associated with tumourigenesis (29), *ESR2* has been shown to inhibit cell proliferation in vitro (30) and tumour formation in nude mice (31). Although we did not identify a somatic *ESR2* alteration in this tumour and were unable to investigate *ESR2* promoter hypermethylation, we note that *ESR2* promoter hypermethylation associated with ER β down-regulation has been detected in a variety of tumour types (32-34). As with other cancer predisposition syndromes (35), it is also probable that additional somatic mutations are required in the transition from normal thyroid to MTC and we did detect low level loss of chromosomes 3 and 13.

It is likely that ER β loss of function in MTC is rare as we did not detect loss of ER β expression in 12 apparently sporadic MTC tumours nor an intragenic *ESR2* mutation in 15 sporadic MTCs. Interestingly, studies of MTC have indicated that *RAS* and *RET* are the dominant driver pathways in tumourigenesis, with few mutations being detected in other genes and none in *ESR2* (36). Our findings are also in keeping with large scale studies of sporadic tumours where no CNVs, and only one putative mutation of *ESR2* (missense alteration in a single papillary thyroid cancer), have been detected in thyroid tumours (18,19). Such relative infrequency of somatic mutation in sporadic tumours in genes predisposing to hereditary cancers is well recognised, for example the low rate of somatic *BRCA1* and *BRCA2* mutations in sporadic breast cancer (37,38). We did note that somatic *ESR2* mutations have been detected in other neuroectodermal derived tumours such as glioma and melanoma and also breast cancer samples indicating that inappropriate up-regulated ER α activity may also be involved in their tumourigenesis (18,19).

We did not detect a constitutional *ESR2* mutation in a further three families with an apparent genetic predisposition to MTC and CCH and it is likely that, akin to other familial cancer predisposition syndromes (39), there is also heterogeneity in the genetic predisposition to MTC/CCH.

The association of increased RET activity with MTC tumourigenesis is well established (4) and here we demonstrate a novel mechanism of MTC tumourigenesis whereby loss of ER β function results in ER α -driven RET expression, likely mediated through the ERE on the *RET* promoter. Whilst there may be an unidentified constitutional *RET* mutation, or another mechanism of *RET* up-regulation, we have also shown *ESR2* c.948delT to cause increased cellular proliferation indicating a likely role in tumourigenesis; however, we cannot exclude that other mechanisms of ER β action may also be involved, for example through loss of ER β mediated inhibition of HIF-1 α causing inappropriate HIF pathway activation (40).

Although we detected a mutation in just one family, our results suggest that constitutional *ESR2* mutation is of high penetrance for CCH/MTC. However, as individual I:1 has not overtly presented with MTC, we cannot exclude that there may be variable penetrance, as occurs for germline *RET* mutations and in other familial cancer predisposition syndromes such as paraganglioma-pheochromocytoma predisposition (4, 41).

It is interesting to speculate whether constitutional *ESR2* mutations are also associated with predisposition to HPT and pheochromocytoma. These tumours have not occurred to date in the family (relevant clinical and laboratory investigation has been undertaken) and whole exome sequencing of both constitutional DNA from individuals with apparent genetic predisposition to pheochromocytoma (12 individuals) and sporadic pheochromocytoma (seven tumours) has not detected any deleterious *ESR2* mutations (ERM unpublished observations). Nonetheless, given the known tumour predisposition associated with dysregulated *RET* activity, it would seem prudent to offer clinical surveillance for pheochromocytoma and HPT, and prophylactic thyroidectomy, where a loss of function constitutional *ESR2* mutation is detected.

The investigation of rare familial cancer predisposition often provides important insights into cell biology with the potential for novel therapeutic approaches (42) and our findings also provide further evidence of the interaction between ER α and RET in tumourigenesis. Many invasive breast

cancers expressing ER α also express RET (4) and in recent pre-clinical models of breast cancer, targeting both RET and ER α decreases the growth and metastatic potential of breast cancer cells (43). Thus tumours associated with loss of ER β function might be suitable candidates for therapeutic agents targeting the RET pathway (44), RET tyrosine kinase active site (45) or ER α (46).

In summary, our data is the first to identify a novel genetic cause for familial MTC since the identification of the *RET* gene. We demonstrate a novel mechanism promoting up-regulation of *RET* activity through constitutional *ESR2* mutation which provides opportunities for pre-symptomatic testing and potential novel therapeutic strategies.

Materials and Methods

Whole Exome Sequencing Analysis

Exome sequencing was performed at the Biomedical Research Centre at Kings College London as previously described (47).

Sanger Sequencing

Genomic DNA was extracted from formalin-fixed paraffin embedded (FFPE) tissue using the QIAamp DNA FFPE kit (Qiagen) and from peripheral leucocytes using the Nucleon BACC2 kit (Amersham Biosciences) (sequences and PCR conditions available upon request).

Immunohistochemistry

Slides were de-paraffinized using Benchmark Special Stains Deparaffinization Solution. Heat-induced antigen retrieval was performed using EDTA buffer as previously described (48). Immunostaining was performed using the Ventana Benchmark Automated Slide Stainer (Ventana Medical Systems) with antibodies against ER β (Leica Biosystems, 1:50), calcitonin (Dako IR515) and RET (Abcam EPR2871, 1:500).

Loss of Heterozygosity (LOH) analyses

Microsatellite markers were used to determine LOH at the *ESR2* locus on chromosome 14q23.2: one, previously described (49) within intron six (chr14:64,720,279-64,720,323) and one 5' upstream (chr14:64,802,692-64,802,742) (sequences available upon request). PCR amplification was performed on genomic DNA templates using fluorescently labelled primers. PCR products were electrophoresed on an ABI Genetic Analyzer and analysed by GeneMapper fragment analysis software (Applied Biosystems).

OncoScan[®] FFPE Assay Kit

The OncoScan[®] FFPE assay kit platform was utilized to identify whole-genome copy number variations (CNV) as previously described (50).

Cell Culture

In the absence of a suitable MTC cell line we used HCT116 (Health Protection Agency Culture Collections, UK) and MCF-7 cells (European Collection of Animal Cell Cultures) cultured in McCoy's 5A (Life Technologies) with 5% fetal bovine serum (Invitrogen), penicillin (10⁴ U/ml) and streptomycin (10⁴ mcg/ml) (Invitrogen) for HCT116 cells and RPMI 1640 (Life Technologies) with 5% fetal bovine serum (Invitrogen), penicillin (10⁵ U/L) and streptomycin (100 mg/L) for MCF-7 cells. Cells were treated with 17 β -oestradiol (E2, Sigma) and 4,4',4''-(4-propyl-[1H]-pyrazole-1,3,5-triyl(trisphenol) (PPT, Tocris) at final concentrations of 10 nM and 100 nM, in phenol red-free medium (Life Technologies) with 5% charcoal-stripped serum.

Transfection

The ESR1/2 ORFS were cloned into SgfI and MluI restriction sites of pCMV6-AN-Myc (Origene). QuikChange II XL site-directed mutagenesis kit (Agilent Technologies) was used to delete the thymine at position 948 of ESR2, generating ESR2, c.948delT. Plasmid DNA transfections were performed with TransIT LT1 (Mirus Bio LLC). In some experiments, plasmid DNA was introduced into cells by reverse transfection (lipid: DNA complexes plated prior to addition of cells).

RNA extraction, Real-time PCR and mRNA Stability Assay

Total RNA was extracted and reverse transcribed using RNeasy Micro Kit (Qiagen) and Reverse Transcription System (Promega) with expression determined by the 7500 Real-time PCR system (Applied Biosystems). qRT-PCR was carried out on 96 well plates using FAM-labelled Taqman probes (Life Technologies) for ESR2 (Hs01100359_m1), RET (Hs01120030_m1) and PPIA (Hs04194521_s1). In mRNA stability experiments, HCT116 cells were treated with actinomycin D at a final concentration of 10 μ g/ml for 4h, 8h and 24h prior to total RNA extraction.

Western Blotting

Western blots, performed as described previously (51), were probed with antibodies against RET (C31B4, Cell Signaling Technology), 1:1000; ER β (H150, Santa Cruz Biotechnology), 1:1000; c-Myc, 1:1000 and β actin; 1:10,000 (Sigma-Aldrich). Antigen-antibody complexes were detected using ECL Plus (Amersham Biosciences).

Luciferase Reporter Assays

Luciferase reporter assays were performed using the Cignal ERE Reporter Kit (Qiagen). HCT116 (7,500 cells/ well) and MCF-7 (15,000 cells/ well) cells were reverse transfected with 50ng Cignal ERE reporter along with 100ng ER expression vectors with TransIT LT1. After 24h hormone incubation, cells were harvested with lysis buffer (Promega) and the firefly and renilla luciferase activities determined with a dual luciferase assay kit (Promega), measuring luminescence with a Centro LB960 microplate luminometer (Berthold Technologies).

Cell Proliferation

Proliferation of MCF-7 cells was determined using the Celltiter-Glo® Luminescent Cell Viability Assay (Promega). In brief, 12,500 MCF-7 cells were reverse transfected with ER expression vectors and TransIT LT1 in 96 wells and then treated with hormones as indicated for 72h prior to analysis of the luminescent signal using the Centro LB960 microplate luminometer (Berthold Technologies).

Statistical Analyses

Data are displayed as mean \pm SE. Normally distributed data were analysed using a two-tailed Student's t-test, unless otherwise indicated. A *P*-value < 0.05 was considered to be statistically significant.

Acknowledgements

This manuscript is dedicated to the memory of Dr. Louise Brueton who undertook all the initial clinical work-up in the family. We are grateful to the families and patients for participation in the study.

The study was funded by the Queen Elizabeth Hospital Birmingham Charity and The Get A-Head Charitable Trust with support from Affymetrix UK Limited, the Technology Strategy Board (now Innovate-UK) Stratified Medicine Innovation Platform (#101032), the Canadian Institutes for Health Research (#142303), the Terry Fox Research Institute Transdisciplinary Training Program in Cancer Research and the National Institute for Health Research.

370

Conflict of Interest

371

372 The authors have no conflicts of interest to declare.

373

References

1. Leboulleux, S., Baudin, E., Travagli, J. P. and Schlumberger, M. (2004) Medullary thyroid carcinoma. *Clin. Endocrinol.*, **61**, 299–310.
2. Modigliani, E., Cohen, R., Campos, J.-M., Conte-Devolx, B., Maes, B., Boneu, A., Schlumberger, M., Bigorgne, J.-C., Dumontier, P., Leclerc, L., *et al.* (1998) Prognostic factors for survival and for biochemical cure in medullary thyroid carcinoma: results in 899 patients. *Clin. Endocrinol.*, **48**, 265–273.
3. Decker, R. A., Peacock, M. L., Borst, M. J., Sweet, J. D. and Thompson, N. W. (1995) Progress in genetic screening of multiple endocrine neoplasia type 2A: is calcitonin testing obsolete? *Surgery*, **118**, 257–264.
4. Mulligan, L. M. (2014) RET revisited: expanding the oncogenic portfolio. *Nat. Rev. Cancer*, **14**, 173–186.
5. Margraf, R.L., Crockett, D. K. and Krautscheid, P. M. F. (2009) Multiple endocrine neoplasia type 2 RET protooncogene database: repository of MEN2-associated RET sequence variation and reference for genotype/phenotype correlations. *Hum. Mutat.*, **30**, 548–556.
6. Wells, S. A., Pacini, F., Robinson, B. G. and Santoro, M. (2013) Multiple endocrine neoplasia type 2 and familial medullary thyroid carcinoma: an update. *J. Clin. Endocrinol. Metab.*, **98**, 3149–3164.
7. Wells, S. A., Asa, S. L., Dralle, H., Elisei, R., Evans, D. B., Gagel, R. F., Lee, N., Machens, A., Moley, J. F., Pacini, F., *et al.* (2015) Revised American Thyroid Association guidelines for the management of medullary thyroid carcinoma. *Thyroid*, **25**, 567–610.
8. Romei, C., Mariotti, S., Fugazzola, L., Taccaliti, A., Pacini, F., Opocher, G., Mian, C., Castellano, M., Uberti, E. Degli, *et al.* (2010) Multiple endocrine neoplasia type 2 syndromes (MEN 2): results from the ItaMEN network analysis on the prevalence of different genotypes and phenotypes, *Eur. J. Endocrinol. Eur. Fed. of Endocr. Soc.*, **163**, 301–308.
9. Fugazzola, L., Cerutti, N., Mannavola, D., Ghilardi, G., Alberti, L., Romoli, R. and Beck-Peccoz, P. (2002) Multigenerational familial medullary thyroid cancer (FMTC): evidence for

402 FMTC phenocopies and association with papillary thyroid cancer, *Clin. Endocrinol.*, **56**, 53–
 403 63.

404 10. Mosselman, S., Polman, J. and Dijkema, R. (1996) ER beta: identification and
 405 characterization of a novel human estrogen receptor. *FEBS Lett.*, **392**, 49–53.

406 11. dbSNP. [Internet]. [cited 2014 Dec 17]. Available from:
 407 http://www.ncbi.nlm.nih.gov/projects/SNP/snp_summary.cgi?build_id=132

408 12. NHLBI Exome Sequencing Project (ESP): Exome Variant Server. [Internet]. [cited 2014 Dec
 409 17]. Available from: <http://evs.gs.washington.edu/EVS/>

410 13. 1000 genomes. [Internet]. [cited 2014 Dec 17]. Available from:
 411 <http://www.1000genomes.org/>

412 14. Rahman, N. (2104) Realizing the promise of cancer predisposition genes. *Nature*, **505**, 302–
 413 308.

414 15. SIFT (J. Craig Venter Institute). [Internet]. [cited 2015 Jun 7]. Available from:
 415 <http://sift.jcvi.org/>

416 16. PolyPhen-2: prediction of functional effects of human nsSNPs. [Internet]. [cited 2015 Jun 7].
 417 Available from: <http://genetics.bwh.harvard.edu/pph2/index.shtml>

418 17. Knudson, A. G. (1971) Mutation and cancer: statistical study of retinoblastoma. *Proc. Natl.*
 419 *Acad. Sci. U S A.*, **68**, 820–823.

420 18. COSMIC: Catalogue of somatic mutations in cancer. [Internet]. [cited 2015 Jun 23].
 421 Available from: <http://cancer.sanger.ac.uk/cosmic>

422 19. The Cancer Genome Atlas (TCGA). [Internet]. [cited 2014 Dec 18]. Available from:
 423 <http://cancergenome.nih.gov/publications/publicationguidelines>

424 20. McKenna, N. J., Xu, J., Nawaz, Z., Tsai, S. Y., Tsai, M. J. and O'Malley, B. W. (1999)
 425 Nuclear receptor coactivators: multiple enzymes, multiple complexes, multiple functions. *J.*
 426 *Steroid. Biochem. Mol. Biol.*, **69**, 3–12.

427 21. Cowley, S. M., Hoare, S., Mosselman, S. and Parker, M. G. (1997) Estrogen receptors alpha
 428 and beta form heterodimers on DNA. *J. Biol. Chem.*, **272**, 19858–19862.

- 429 22. Hall, J. M. and McDonnell, D. P. (1999) The estrogen receptor beta-isoform (ERbeta) of the
430 human estrogen receptor modulates ERalpha transcriptional activity and is a key regulator of
431 the cellular response to estrogens and antiestrogens. *Endocrinology*, **140**, 5566–5578.
- 432 23. Pace, P., Taylor, J., Suntharalingam, S., Coombes, R. C. and Ali, S. (1997) Human estrogen
433 receptor beta binds DNA in a manner similar to and dimerizes with estrogen receptor alpha. *J.*
434 *Biol. Chem.*, **272**, 25832–25838.
- 435 24. Morandi, A., Plaza-Menacho, I. and Isacke, C. M. (2011) RET in breast cancer: functional
436 and therapeutic implications. *Trends. Mol. Med.*, **17**, 149–157.
- 437 25. Boulay, A., Breuleux, M., Stephan, C., Fux, C., Briskin, C., Fiche, M., Wartmann, M.,
438 Stumm, M., Lane, H. A. and Hynes, N. E. (2008) The Ret Receptor Tyrosine Kinase Pathway
439 Functionally Interacts with the ERα Pathway in Breast Cancer. *Cancer. Res.*, **68**, 3743–3751.
- 440 26. Frasor, J., Stossi, F., Danes, J. M., Komm, B., Lyttle, C. R. and Katzenellenbogen, B. S.
441 (2004) Selective estrogen receptor modulators: discrimination of agonistic versus antagonistic
442 activities by gene expression profiling in breast cancer cells. *Cancer Res.*, **64**, 1522–1533.
- 443 27. Tozlu, S., Girault, I., Vacher, S., Vendrell, J., Andrieu, C., Spyrtos, F., Cohen, P., Lidereau,
444 R. and Bieche, I. (2006) Identification of novel genes that co-cluster with estrogen receptor
445 alpha in breast tumor biopsy specimens, using a large-scale real-time reverse transcription-
446 PCR approach. *Endocr. Relat. Cancer*, **13**, 1109–1120.
- 447 28. Taylor, A. H. and Al-Azzawi, F. (2000) Immunolocalisation of oestrogen receptor beta in
448 human tissues. *J. Mol. Endocrinol.*, **24**, 145–155.
- 449 29. Kregge, J.H., Hodgins, J.B., Couse, J.F., Enmark, E., Warner, M., Mahler, J.F., Sar, M., Korach,
450 K.S., Gustafsson, J.A. and Smithies, O. (1998) Generation and reproductive fitness of mice
451 lacking estrogen receptor beta. *Proc. Natl. Acad. Sci.*, **95**, 15677–15682.
- 452 30. Lazennec, G., Bresson, D., Lucas, A., Chauveau, C. and Vignon, F. (2001) ER beta inhibits
453 proliferation and invasion of breast cancer cells. *Endocrinology*, **142**, 4120–4130.
- 454 31. Paruthiyil, S., Parmar, H., Kerekatte, V., Cunha, G. R., Firestone, G. L. and Leitman, D. C.
455 (2004) Estrogen receptor beta inhibits human breast cancer cell proliferation and tumor
456 formation by causing a G2 cell cycle arrest. *Cancer Res.*, **64**, 423–428.

32. Al-Nakhle, H., Smith, L., Bell, S. M., Burns, P. A., Cummings, M., Hanby, A. M., Lane, S., Parker, M. D., Hughes, T. A. and Speirs, V. (2013) Regulation of estrogen receptor β 1 expression in breast cancer by epigenetic modification of the 5' regulatory region. *Int. J. Oncol.*, **43**, 2039–2045.
33. Suzuki, F., Akahira, J., Miura, I., Suzuki, T., Ito, K., Hayashi, S., Sasano, H. and Yaegashi, N. (2008) Loss of estrogen receptor β isoform expression and its correlation with aberrant DNA methylation of the 5'-untranslated region in human epithelial ovarian carcinoma. *Cancer Sci.*, **99**, 2365–2372.
- 34.. Zhu, X., Leav, I., Leung, Y.-K., Wu, M., Liu, Q., Gao, Y., McNeal, J. E. and Ho, S.-M. (2004) Dynamic regulation of estrogen receptor-beta expression by DNA methylation during prostate cancer development and metastasis. *Am. J. Path.*, **164**, 2003–2012.
35. Fearon, E. R. and Vogelstein, B. (1990) A genetic model for colorectal tumorigenesis. *Cell*, **61**, 759–767.
36. Agrawal, N., Jiao, Y., Sausen, M., Leary, R., Bettegowda, C., Roberts, N. J., Bhan, S., Ho, A. S., Khan, Z., Bishop, J., *et al.* (2013) Exomic Sequencing of Medullary Thyroid Cancer Reveals Dominant and Mutually Exclusive Oncogenic Mutations in RET and RAS. *J. Clin. Endocrinol. Metab.*, **98**, E364–E369.
37. Futreal, P. A., Liu, Q., Shattuck-Eidens, D., Cochran, C., Harshman, K., Tavtigian, S., Bennett, L. M., Haugen-Strano, A., Swensen, J. and Miki, Y. (1994) BRCA1 mutations in primary breast and ovarian carcinomas. *Science*, **266**, 120–122.
38. Lancaster, J. M., Wooster, R., Mangion, J., Phelan, C. M., Cochran, C., Gumbs, C., Seal, S., Barfoot, R., Collins, N., Bignell, G., *et al.* (1996) BRCA2 mutations in primary breast and ovarian cancers. *Nat. Genet.*, **13**, 238–240.
39. Stratton, M.R. and Rahman, N. (2008) The emerging landscape of breast cancer susceptibility. *Nat Genet.*, **40**, 17–22.
40. Lim, W., Park, Y., Cho, J., Park, C., Park, J., Park, Y.-K., Park, H. and Lee, Y. (2011) Estrogen receptor beta inhibits transcriptional activity of hypoxia inducible factor-1 through

the downregulation of arylhydrocarbon receptor nuclear translocator, *Breast Cancer Res.*, **13**, R32.

41. Maher, E.R. (2014) Pheochromocytoma and paraganglioma: next-generation sequencing and evolving Mendelian syndromes. *Clin. Med.*, **14**, 440–444.
42. Gossage, L., Eisen, T. and Maher, E. R. (2015) VHL, the story of a tumour suppressor gene. *Nat. Rev. Cancer.*, **15**, 55-64.
43. Spanheimer, P. M., Park, J.-M., Askeland, R. W., Kulak, M. V., Woodfield, G. W., De Andrade, J. P., Cyr, A. R., Sugg, S. L., Thomas A. and Weigel, R. J. (2014) Inhibition of RET increases the efficacy of antiestrogen and is a novel treatment strategy for luminal breast cancer. *Clin. Cancer Res.*, **20**, 2115–2125.
44. Wells, S. A., Robinson, B. G., Gagel, R. F., Dralle, H., Fagin, J. A., Santoro, M., Baudin, E., Elisei, R., Jarzab, B., Vasselli, J. R., *et al.* (2012) Vandetanib in Patients With Locally Advanced or Metastatic Medullary Thyroid Cancer: A Randomized, Double-Blind Phase III Trial. *J. Clin. Oncol.*, **30**, 134–141.
45. De Falco, V., Buonocore, P., Muthu, M., Torregrossa, L., Basolo, F., Billaud, M., Gozgit, J. M., Carlomango, F. and Santoro, M. (2013) Ponatinib (AP24534) is a novel potent inhibitor of oncogenic RET mutants associated with thyroid cancer. *J. Clin. Endocrinol. Metab.*, **98**, E811–E819.
46. Jordan, V. C. and Koerner, S. (1975) Tamoxifen (ICI 46,474) and the human carcinoma 8S oestrogen receptor. *Eur. J. Cancer*, **11**, 205-206.
47. Jafri, M., Wake, N. C., Ascher, D. B., Pires, D. E. V., Gentle, D., Morris, M. R., Rattenberry, E., Simpson, M. A., Trembath, R. C., Weber, A., *et al.* (2015) Germline Mutations in the CDKN2B Tumor Suppressor Gene Predispose to Renal Cell Carcinoma. *Cancer Discov.*, **5**, 723–729.
48. Myers, J. (2004) Automated slide stainers for special stains, immunohistochemistry and in situ hybridization: A review. *Med. Lab. Observer*, **36**, 28-30.

49. Tsukamoto, K., Inoue, S., Hosoi, T., Orimo, H. and Emi, M. (1998) Isolation and radiation hybrid mapping of dinucleotide repeat polymorphism at the human estrogen receptor beta locus. *J. Hum. Genet.*, **43**, 73–74.
50. Foster, J. M., Oumie, A., Togneri, F. S., Vasques, F. R., Hau, D., Taylor, M., Tinkler-Hundal, E., Southward, K., Medlow, P., McGreeghan-Crosby, K., *et al.* (2015) Cross-laboratory validation of the OncoScan® FFPE Assay, a multiplex tool for whole genome tumour profiling. *BMC Med. Genomics*, **8**, 5.
51. Smith, V. E., Read, M. L., Turnell, A. S., Watkins, R. J., Watkinson, J. C., Lewy, G. D., Fong, J. C. W., James, S. R., Eggo, M. C., Boelaert, K., *et al.* (2009) A novel mechanism of sodium iodide symporter repression in differentiated thyroid cancer. *J. Cell. Sci.*, **122**, 3393–3402.

Figure legends

Figure 1. Germline *ESR2* mutation detected in family members with MTC/CCH and loss of ER β staining in associated MTC tumour. (A) Sanger sequencing traces showing presence of *ESR2* c.948delT (upper panel) and corresponding normal trace (lower panel). Affected nucleotide is indicated by a red arrow. (B) Representative histological stains in sections following thyroid surgery in representative family members II:3 (MTC), III:2 (CCH), III:4 (normal thyroid). Haematoxylin and eosin stain showing MTC (upper panel) and apparently normal thyroid tissue (middle and lower panels). Calcitonin staining demonstrating MTC (upper panel), areas of CCH within apparently normal thyroid tissue (black arrowhead, middle panel), and normal thyroid tissue (lower panel). ER β immunohistochemistry showing loss of nuclear expression in the MTC (upper panel, white arrowhead). Middle and lower panels – preserved nuclear staining of ER β in follicular epithelial cells. It was not possible to determine if there was loss of ER β staining associated with areas of CCH (middle panel).

Figure 2. Germline *ESR2* alteration detected in an individual with early onset MTC and preserved ER β staining in associated MTC tumour. (A) Sanger sequencing traces showing constitutional *ESR2* alteration c.382G>C (upper panel) and corresponding normal trace (lower panel) detected in an individual with apparently isolated MTC age 36 years. Affected nucleotide is indicated by a red arrow. (B) Representative histological stains in sections following thyroid surgery. Left, haematoxylin and eosin stain showing MTC (white arrowhead) and corresponding normal thyroid tissue (black arrowhead). Middle, calcitonin stain demonstrating MTC (white arrowhead) with corresponding unstained normal thyroid tissue (black arrowhead). Right, ER β immunohistochemistry showing preserved nuclear expression in both MTC (white arrowhead and high power insert) and corresponding normal thyroid tissue (black arrowhead and high power insert).

Figure 3. *ESR2* c.948delT is associated with unrestrained ER α -driven cellular proliferation. MCF-7 cells were reverse transfected with wtESR1 (ESR1) alone or in combination with either wt*ESR2* (ESR2) or *ESR2* c.948delT (948 Δ T) for 24h and then treated with 10nM E2, 100nM E2, or 10nM PPT

for 72h, compared with vehicle (ethanol) treated controls. Cell proliferation status was quantified by luciferase assay using the CellTiter-Glo Luminescent Cell Viability Assay (Promega) and expressed relative to ethanol treated controls. Data presented as mean values \pm SE (n = 4). *, P < 0.05; **, P < 0.01; ***, P < 0.001; NS - not significant.

Figure 4. Diminished ability of *ESR2* c.948delT to inhibit wt*ESR1*. Luciferase assays evaluating transcriptional activity of a transfected ERE by vector only (VO), wt*ESR1* (ESR1) alone, or in combination with either wt*ESR2* (ESR2) or *ESR2* c.948delT (948 Δ T) mutant at a ratio of 1:3 in (A) HCT116 and (B) MCF-7 cell lines. Cells were treated with 10nM E2, 10nM PPT or ethanol control as indicated for 24h prior to cell lysis and luciferase activity measurement. Relative fold change presents mean values \pm SE from at least two independent experiments (n = 4 wells per experiment). *, P < 0.05; **, P < 0.01; ***, P < 0.001. NS - not significant.

Figure 5. *ESR2* c.948delT is associated with elevated RET expression. (A,B) real-time PCR (upper panel) and Western blot (lower panel) analyses of RET expression levels in HCT116 cells transfected with wt*ESR1* (ESR1) alone or in combination with either wt*ESR2* (ESR2) or *ESR2* c.948delT (948 Δ T) and then treated with either (A) 10nM E2 (+) or (B) 10nM PPT (+) compared to ethanol control (-). Real-time data presented as mean *RET* mRNA levels \pm SE (n = 4). In control experiments cells were transfected with vector only (VO) as indicated. (C) Representative histological sections showing immunohistochemical staining for RET (upper panels) with corresponding calcitonin stains (lower panels) to confirm presence of CCH/MTC. Left - archived slides from individual II:1 showing areas of intense RET staining (white arrow, upper panel) and area of marked calcitonin staining indicating presence of CCH (white arrowhead, lower panel) within normal thyroid tissue. Middle - MTC tumour from individual II:3 showing intense RET expression (white arrowhead, upper panel) as compared with corresponding normal thyroid tissue (black arrowhead, upper panel). Calcitonin stain (lower panel) confirms presence of MTC (same panel as in Fig.1B). Right - MEN2 associated MTC (white arrowhead, upper panel) also showing intense RET expression as compared with corresponding normal thyroid tissue (black arrowhead, upper panel), for comparison with middle

579 panel. Calcitonin stain (lower panel) confirms presence of MTC (white arrowhead, lower panel) with
580 corresponding normal thyroid tissue also shown (black arrowhead, lower panel).
581

Abbreviations

582		
583		
584	MTC	medullary thyroid cancer
585	CCH	C cell hyperplasia
586	ER β	oestrogen receptor beta
587	ERE	estrogen response element
588	MEN2	multiple endocrine neoplasia type 2
589	HPT	hyperparathyroidism
590	SNP	single nucleotide polymorphism
591	NHLBI	National Heart, Lung and Blood Institute
592	EVS	exome variant server
593	CNV	copy number variant
594	SIFT	Sorts Intolerant From Tolerant
595	TCGA	The Cancer Genome Atlas
596	ER α	oestrogen receptor alpha
597	E2	17 β -estradiol
598	PPT	4,4',4''-(4-propyl-[1H]-pyrazole-1,3,5-triyl(trisphenol)
599	FFPE	formalin-fixed paraffin embedded
600	PCR	polymerase chain reaction
601	LOH	loss of heterozygosity

602 COSMIC catalogue of somatic mutations in cancer

603

Figure 1





Figure 2

A



B

Haematoxylin-eosin

Calcitonin

ER β








Figure 3




Figure 4





Figure 5



SUPPLEMENTARY FIGURES




Figure S1. Pedigree of family with MTC/CCH. Diagnoses marked are from histological review following thyroid surgery. The index case (arrowed) presented with MTC age 22 years. ESR2 c.948delT carriers are marked and the ages (y) at which thyroidectomy was undertaken are indicated.




Figure S2. OncoScan® FFPE assay of MTC tumour. Log2 ratio (upper panel) and BAF plots (lower panel) showing results from the OncoScan® FFPE Assay Kit for MTC tumour from individual II:3. Data shows no copy number abnormality of either chromosome 10 or 14. There is some evidence of low level loss of chromosomes 3 and 13.




Figure S3. ESR2 c.948delT results in reduced mRNA levels. At 24 hours after transfection with either wt ESR2 (ESR2, white diamonds) or ESR2 c.948delT (948ΔT, black squares), time 0 hours (h) on x-axis, HCT116 cells were treated with actinomycin D (10 mg/ml) for 0, 4, 8 and 24 hours. The relative wt ESR2 and c.948delT mRNA levels were monitored by RT-qPCR. All values are relative to wt ESR2 mRNA levels at 0 hours (set as 1). Values represent means \pm SE (n = 4) from at least two independent experiments. ESR2 mRNA expression levels were normalized against the 18S rRNA housekeeping gene. **, $P < 0.01$; ***, $P < 0.001$. NS - not significant.

<i>AIP</i>	<i>DDB2</i>	<i>FANCL</i>	<i>PALB2</i>	<i>SLX4</i>
<i>ALK</i>	<i>DICER1</i>	<i>FANCM</i>	<i>PHOX2B</i>	<i>SMAD4</i>
<i>APC</i>	<i>DIS3L2</i>	<i>FH</i>	<i>PMS1</i>	<i>SMARCB1</i>
<i>ATM</i>	<i>EGFR</i>	<i>FLCN</i>	<i>PMS2</i>	<i>STK11</i>
<i>BAP1</i>	<i>EPCAM</i>	<i>GATA2</i>	<i>PRF1</i>	<i>SUFU</i>
<i>BLM</i>	<i>ERCC2</i>	<i>GPC3</i>	<i>PRKAR1A</i>	<i>TMEM127</i>
<i>BMPR1A</i>	<i>ERCC3</i>	<i>HNF1A</i>	<i>PTCH1</i>	<i>TP53</i>
<i>BRCA1</i>	<i>ERCC4</i>	<i>HRAS</i>	<i>PTEN</i>	<i>TSC1</i>
<i>BRCA2</i>	<i>ERCC5</i>	<i>KIT</i>	<i>RAD51C</i>	<i>TSC2</i>
<i>BRIP1</i>	<i>EXT1</i>	<i>MAX</i>	<i>RAD51D</i>	<i>VHL</i>
<i>BUB1B</i>	<i>EXT2</i>	<i>MEN1</i>	<i>RB1</i>	<i>WRN</i>
<i>CDC73</i>	<i>EZH2</i>	<i>MET</i>	<i>RECQL4</i>	<i>WT1</i>
<i>CDH1</i>	<i>FANCA</i>	<i>MLH1</i>	<i>RET</i>	<i>XPA</i>
<i>CDK4</i>	<i>FANCB</i>	<i>MSH2</i>	<i>RHBDF2</i>	<i>XPC</i>
<i>CDKN1C</i>	<i>FANCC</i>	<i>MSH6</i>	<i>RUNX1</i>	
<i>CDKN2A</i>	<i>FANCD2</i>	<i>MUTYH</i>	<i>SBDS</i>	
<i>CEBPA</i>	<i>FANCE</i>	<i>NBN</i>	<i>SDHAF2</i>	
<i>CEP57</i>	<i>FANCF</i>	<i>NF1</i>	<i>SDHB</i>	
<i>CHEK2</i>	<i>FANCG</i>	<i>NF2</i>	<i>SDHC</i>	
<i>CYLD</i>	<i>FANCI</i>	<i>NSD1</i>	<i>SDHD</i>	

Table S1. List of genes on TruSight Cancer Panel.

(http://www.illumina.com/products/trusight_cancer.html)